

NEW MEXICO CROP PROGRESS



United States Department of Agriculture
NATIONAL AGRICULTURAL STATISTICS SERVICE
NEW MEXICO FIELD OFFICE
 PO Box 1809, Las Cruces, NM 88004
 Cooperating with the New Mexico Department of Agriculture



FOR IMMEDIATE RELEASE
 June 26, 2017

Contact: Longino Bustillos
 (800) 530-8810

CROP PROGRESS AND CONDITION WEEK ENDING JUNE 25, 2017

AGRICULTURAL SUMMARY: With year-to-date precipitation totals at or below normal at 20 of the 45 reporting weather stations and totals at the remaining 25 falling quickly, scorching temperatures continued to dominate the State during the week, according to the Mountain Regional Field Office of the National Agricultural Statistical Service, USDA. While measurable moisture was recorded in most northeastern and southern locations, rainfall totals were short when compared with average, and did little to help alleviate the prolonged dryness evident in some areas. Comments from Grant and Luna Counties indicated that moisture was widespread; however, it immediately soaked into dry soils, and without a considerable amount of precipitation over the next few months, pastures will not provide adequate feed during the winter. Lake Roberts, at 1.37 inches, reported the largest moisture accumulation during the week. Triple-digit temperatures were recorded at 29 of the 45 reporting weather stations, with daytime highs ranging from 81 degrees at Angel Fire to 111 degrees at Acme. Reports from Dona Ana County indicated that milk production at several dairies declined as a result of the extremely hot conditions. Average temperatures ranged from normal to 11 degrees above average. Overnight lows varied from 34 degrees at Ocate to 69 degrees at Carlsbad. Statewide, topsoil moisture levels were reported as 37 percent adequate to surplus, compared with 38 percent last week, 28 percent last year, and a 5-year average of 23 percent. As the onion harvest surpassed the halfway point during the week, 84 percent of the crop was reported in good to excellent condition, compared with 92 percent last year. Hail ranging from pea-size to approximately 1 inch in diameter was noted in portions of the northeast. Hail damage in all crops was reported as 5 percent light, compared with 1 percent light last week. Wind damage in all crops was reported as 3 percent moderate and 14 percent light, compared with 2 percent moderate and 23 percent light last week. Stock water supplies were reported as 14 percent very short, 34 percent short, and 52 percent adequate, compared with 9 percent very short, 31 percent short, and 60 percent adequate last week.

CROP AND LIVESTOCK PROGRESS

Commodity	Current week (percent)	Previous week (percent)	Previous year (percent)	5-year average (percent)
Alfalfa hay				
1 st cutting harvested.....	96	94	96	98
2 nd cutting harvested.....	56	46	57	66
3 rd cutting harvested.....	24	13	3	19
Corn				
Emerged.....	93	76	81	84
Cotton				
Planted.....	96	93	95	99
Emerged.....	83	75	87	NA
Squaring.....	7	--	9	23
Onions				
Harvested.....	55	48	39	56
Peanuts				
Planted.....	93	85	86	96
Emerged.....	82	73	60	NA
Sorghum				
Planted.....	73	64	82	78
Emerged.....	32	27	54	NA
Winter wheat				
Harvested for grain.....	77	48	61	49

NA – not available
 (--) – zero

DAYS SUITABLE FOR FIELDWORK AND SOIL MOISTURE CONDITION

Commodity	Current week	Previous week	Previous year	5-year average
Days suitable for fieldwork.....	6.6	7.0	6.7	6.4
Topsoil moisture	(percent)	(percent)	(percent)	(percent)
Very short.....	22	16	12	44
Short.....	41	46	60	33
Adequate.....	36	37	27	23
Surplus.....	1	1	1	--
Subsoil moisture				
Very short.....	18	13	10	NA
Short.....	51	49	40	NA
Adequate.....	30	37	50	NA
Surplus.....	1	1	--	NA

NA – not available
 (--) – zero

CROP, LIVESTOCK, AND PASTURE AND RANGE CONDITION

	Current week	Previous week	Previous year	5-year average
	(percent)	(percent)	(percent)	(percent)
Alfalfa hay				
Very poor	--	--	--	1
Poor.....	1	9	4	5
Fair.....	26	22	46	32
Good	69	68	42	48
Excellent	4	1	8	14
Chile				
Very poor	--	--	3	NA
Poor.....	3	2	12	NA
Fair.....	54	55	16	NA
Good	40	35	48	NA
Excellent	3	8	21	NA
Corn				
Very poor	1	1	--	2
Poor.....	3	1	--	3
Fair.....	40	41	31	33
Good	38	30	51	40
Excellent	18	27	18	22
Cotton				
Very poor	--	--	2	2
Poor.....	7	6	27	13
Fair.....	59	53	29	36
Good	32	35	33	31
Excellent	2	6	9	18
Onion				
Very poor	--	--	--	NA
Poor.....	--	--	--	NA
Fair.....	16	16	8	NA
Good	69	55	45	NA
Excellent	15	29	47	NA
Pasture and range condition				
Very poor	7	4	2	32
Poor.....	24	29	18	26
Fair.....	45	40	48	24
Good	19	23	29	15
Excellent	5	4	3	3
Peanuts				
Very poor	--	--	--	NA
Poor.....	2	--	--	NA
Fair.....	61	61	100	NA
Good	37	39	--	NA
Excellent	--	--	--	NA
Pecan				
Very poor	--	--	--	--
Poor.....	--	--	--	1
Fair.....	4	3	9	19
Good	71	80	54	65
Excellent	25	17	37	15
Sorghum				
Very poor	--	--	--	15
Poor.....	7	9	--	10
Fair.....	58	60	72	34
Good	35	31	28	40
Excellent	--	--	--	1
Winter wheat				
Very poor	10	7	2	NA
Poor.....	15	14	16	NA
Fair.....	31	35	35	NA
Good	26	32	42	NA
Excellent	18	12	5	NA
Cattle and calves				
Very poor	1	--	2	NA
Poor.....	4	1	3	NA
Fair.....	45	39	33	NA
Good	46	52	54	NA
Excellent	4	8	8	NA
Sheep and lambs				
Very poor	3	6	12	NA
Poor.....	8	9	11	NA
Fair.....	25	16	18	NA
Good	64	69	57	NA
Excellent	--	--	2	NA

NA – not available

(--) – zero

New Mexico’s weather data can be accessed at the following:

[http://www.nass.usda.gov/Statistics by State/New Mexico/Publications/Crop Progress & Condition/2017/NM Weather 06252017.pdf](http://www.nass.usda.gov/Statistics_by_State/New_Mexico/Publications/Crop_Progress_&_Condition/2017/NM_Weather_06252017.pdf)