

NEW MEXICO CROP PROGRESS



United States Department of Agriculture
NATIONAL AGRICULTURAL STATISTICS SERVICE
NEW MEXICO FIELD OFFICE
 PO Box 1809, Las Cruces, NM 88004
 Cooperating with the New Mexico Department of Agriculture



FOR IMMEDIATE RELEASE
 September 11, 2017

Contact: Longino Bustillos
 (800) 530-8810

CROP PROGRESS AND CONDITION WEEK ENDING SEPTEMBER 10, 2017

AGRICULTURAL SUMMARY: Scattered showers were experienced around most of the State last week however weekly totals of accumulation were well below normal for all but two weather stations, according to the Mountain Regional Field Office of the National Agricultural Statistical Service, USDA. Temperatures were also above normal for 35 out of the 45 weather stations. The warm weather and lack of moisture allowed for 7 days suitable for fieldwork, compared with the five year average of 6.2 days. Lake Roberts weather station received the largest amount of rainfall for the week at 0.63 inches. Comments from several counties stated the warm weather helped with the maturing of row crops such as corn and sorghum and allowed for winter wheat planting. Statewide, topsoil moisture levels were reported as 70 percent adequate to surplus, compared with 77 percent last week, 42 percent last year, and a 5-year average of 43 percent. A continued decline in pasture and range conditions was experienced due to lack of sufficient moisture, with 36 percent reported in good to excellent condition, compared with 40 percent last week, 30 percent last year, and a 5-year average of 28 percent. Average temperatures ranged from 3 degrees below to 8 degrees above normal. Daytime highs varied from 77 degrees at Angel Fire to 96 degrees at Hachita. Overnight lows ranged from 32 degrees at Ocate to 63 degrees at Santa Teresa and Truth or Consequences. Pecan nut set was reported as 64 percent moderate and 36 percent light, compared with 53 percent moderate, and 47 percent light last week. Hail damage in all crops was reported as 6 percent severe, 7 percent moderate, and 12 percent light, compared with 6 percent severe, 8 percent moderate, and 9 percent light last week. Wind damage in all crops was reported as 7 percent severe, 7 percent moderate, and 29 percent light, compared with 5 percent severe, 6 percent moderate, and 18 percent light last week. Stock water supplies were reported as 4 percent very short, 19 percent short, 75 percent adequate, and 2 percent surplus, compared with 3 percent very short, 17 percent short, 77 percent adequate, and 3 percent surplus last week.

CROP AND LIVESTOCK PROGRESS

| Commodity | Current week (percent) | Previous week (percent) | Previous year (percent) | 5-year average (percent) |
|--|---------------------------|----------------------------|----------------------------|-----------------------------|
| Alfalfa hay | | | | |
| 4 th cutting harvested..... | 85 | 78 | 90 | 96 |
| 5 th cutting harvested..... | 46 | 42 | 53 | 62 |
| 6 th cutting harvested..... | 17 | 8 | 2 | 5 |
| Chile | | | | |
| Green harvested..... | 70 | 64 | 77 | 73 |
| Corn | | | | |
| Dough..... | 70 | 62 | 84 | 88 |
| Dented..... | 44 | 32 | 48 | 54 |
| Mature..... | 15 | 1 | 3 | 14 |
| Corn harvested for silage..... | 50 | 33 | 43 | 48 |
| Cotton | | | | |
| Setting bolls..... | 85 | 81 | 95 | 93 |
| Bolls opening..... | 12 | 9 | 57 | 34 |
| Sorghum | | | | |
| Headed..... | 81 | 72 | 91 | 86 |
| Coloring..... | 72 | 53 | 43 | 26 |
| Mature..... | 4 | 2 | 9 | 2 |
| Winter wheat | | | | |
| Planted..... | 48 | 23 | 15 | 15 |
| Emerged..... | 4 | NA | NA | NA |

NA – not available
 (--) – zero

DAYS SUITABLE FOR FIELDWORK AND SOIL MOISTURE CONDITION

| Commodity | Current week | Previous week | Previous year | 5-year average |
|----------------------------------|--------------|---------------|---------------|----------------|
| Days suitable for fieldwork..... | 7.0 | 7.0 | 6.4 | 6.2 |
| Topsoil moisture | (percent) | (percent) | (percent) | (percent) |
| Very short..... | 5 | 4 | 8 | 24 |
| Short..... | 25 | 19 | 50 | 33 |
| Adequate..... | 64 | 68 | 37 | 38 |
| Surplus..... | 6 | 9 | 5 | 5 |
| Subsoil moisture | | | | |
| Very short..... | 3 | 3 | 8 | NA |
| Short..... | 30 | 22 | 30 | NA |
| Adequate..... | 61 | 65 | 60 | NA |
| Surplus..... | 6 | 10 | 2 | NA |

NA – not available
 (--) – zero

CROP, LIVESTOCK, AND PASTURE AND RANGE CONDITION

| | Current week (percent) | Previous week (percent) | Previous year (percent) | 5-year average (percent) |
|--------------------------|---------------------------|----------------------------|----------------------------|-----------------------------|
| Alfalfa hay | | | | |
| Very poor..... | -- | 1 | -- | 2 |
| Poor..... | 25 | 17 | 4 | 6 |
| Fair..... | 20 | 14 | 42 | 25 |
| Good..... | 45 | 51 | 45 | 57 |
| Excellent..... | 10 | 17 | 9 | 10 |
| Corn | | | | |
| Very poor..... | 5 | 5 | 2 | 2 |
| Poor..... | 6 | 6 | 3 | 5 |
| Fair..... | 25 | 27 | 31 | 23 |
| Good..... | 56 | 47 | 46 | 40 |
| Excellent..... | 8 | 15 | 18 | 30 |
| Cotton | | | | |
| Very poor..... | 5 | 3 | 3 | 2 |
| Poor..... | 7 | 4 | 25 | 13 |
| Fair..... | 28 | 35 | 32 | 34 |
| Good..... | 48 | 40 | 33 | 32 |
| Excellent..... | 12 | 18 | 7 | 19 |
| Pasture and range | | | | |
| Very poor..... | 1 | 1 | 2 | 18 |
| Poor..... | 22 | 22 | 22 | 22 |
| Fair..... | 41 | 37 | 46 | 32 |
| Good..... | 29 | 34 | 26 | 23 |
| Excellent..... | 7 | 6 | 4 | 5 |
| Peanuts | | | | |
| Very poor..... | -- | -- | -- | 5 |
| Poor..... | 8 | 4 | 3 | 17 |
| Fair..... | 61 | 66 | 69 | 64 |
| Good..... | 31 | 30 | 28 | 13 |
| Excellent..... | -- | -- | -- | 1 |
| Pecan | | | | |
| Very poor..... | -- | -- | -- | -- |
| Poor..... | 1 | 3 | -- | -- |
| Fair..... | 6 | 3 | 9 | 14 |
| Good..... | 69 | 70 | 53 | 58 |
| Excellent..... | 24 | 24 | 38 | 28 |
| Sorghum | | | | |
| Very poor..... | 9 | 8 | -- | 9 |
| Poor..... | 2 | 1 | 2 | 7 |
| Fair..... | 39 | 48 | 75 | 36 |
| Good..... | 45 | 38 | 22 | 43 |
| Excellent..... | 5 | 5 | 1 | 5 |
| Cattle and calves | | | | |
| Very poor..... | -- | -- | 2 | NA |
| Poor..... | 1 | 1 | 4 | NA |
| Fair..... | 29 | 36 | 36 | NA |
| Good..... | 63 | 56 | 53 | NA |
| Excellent..... | 7 | 7 | 5 | NA |
| Sheep and lambs | | | | |
| Very poor..... | 2 | 1 | 12 | NA |
| Poor..... | 4 | 2 | 12 | NA |
| Fair..... | 27 | 22 | 16 | NA |
| Good..... | 64 | 65 | 54 | NA |
| Excellent..... | 3 | 10 | 6 | NA |

NA – not available

(--) – zero

New Mexico’s weather data can be accessed at the following:

http://www.nass.usda.gov/Statistics_by_State/New_Mexico/Publications/Crop_Progress_&_Condition/2017/NM_Weather_09102017.pdf